

NOTES ON NEW AND NOTEWORTHY PLANTS. CLXXIX

Harold N. Moldenke

VERBENA PARVULA f. *ALBIFLORA* Mold., f. nov.

Haec forma a forma speciei corollis albis recedit.

This form differs from the typical form of the species in having white corollas.

The form is based on *E. Bastian* 36, collected in "suelo franco arenoso, pH 7, S.-Exp. 3^o, area del Ceramitar, Prov. Cercado", at about 1900 meters altitude, in Tarija, Bolivia, on December 4, 1985, deposited in the Lundell Herbarium at the University of Texas

PAEPALANTHUS AMOENUS var. *BOLIVIANUS* Mold., var. nov.

Haec varietas a forma typica speciei differt inflorescentiis paucicapitulatis numerosis in parte superiori caulis dispersis, nodis caulinis prominente incrassatis.

This variety differs from the typical form of the species chiefly in having its inflorescences terminating abbreviated branches along the upper part of the erect stem, the stems prominently enlarged at the points of emergence of the short and slender branches, the branches densely covered with antrorse basally appressed leaves, the terminal umbels comprising 2--4 unicapitate peduncles, the sheaths about 3 cm. long, the peduncles about 20 cm. long, and the globose many-flowered heads about 1 cm. long and wide.

The variety is based on *R. Haase* 685 from Palmar, on a wet savanna west of the Rio Beni, at 180 meters altitude, Luisita, 13°5' S, 67°15' W, in the province of Iturralde, La Paz, Bolivia, collected on September 2, 1985, and deposited in the Lundell Herbarium at the University of Texas, Austin.